

Ability to kneel after total knee replacement

S. H. Palmer, C. T. Servant, J. Maguire, E. N. Parish, M. J. Cross

From the Australian Institute of Musculo-Skeletal Research, Crows Nest, Australia

We have investigated the ability to kneel after total knee replacement. We asked 75 patients (100 knees) at least six months after routine uncemented primary total knee replacement, to comment on and to demonstrate their ability to kneel. Differences between the perceived and actual ability to kneel were noted. In 32 knees patients stated that they could kneel without significant discomfort. In 54 knees patients avoided kneeling because of uncertainties or recommendations from third parties (doctors, nursing staff, friends, etc). A total of 64 patients was actually able to kneel without discomfort or with mild discomfort only and 12 of the remainder were unable to kneel because of problems which were not related to the knee. Twenty-four patients therefore were unable to kneel because of discomfort in the knee.

There was no difference between the 'kneelers' and 'non-kneelers' with regard to overall knee score, range of movement and the presence of patellar resurfacing.

J Bone Joint Surg [Br] 2002;84-B:220-2.

Received 27 June 2001; Accepted after revision 3 August 2001

There have been a number of studies of knee replacement systems which have shown adequate functional results at middle- to long-term follow-up.¹⁻⁶ Most of the functional scoring systems quoted in these studies use pain, the ability to walk or to ascend and descend stairs, the use of a walking aid, etc as measurements of outcome.⁷⁻¹⁰ The ability to kneel is usually not considered. Some recently

introduced scoring systems, derived from patient questionnaires, include kneeling as a criterion of function of the knee.^{7,11,12}

The kneeling position is important in many activities of daily living and in certain occupations.^{13,14} It has been shown to be a predisposing factor for osteoarthritis of the knee and so preselects patients who will need to kneel after surgery. Kneeling has also been shown to be an intermediate position used by older adults to enable them to rise from the floor.¹⁵

Many patients advised to have surgery for arthritis of the knee enquire about the ability to kneel after operation, but there is little published information on the subject. We have therefore assessed subjectively and objectively the ability of patients to kneel after total knee replacement.

Patients and Methods

We studied 100 knees in 75 consecutive patients attending a follow-up clinic after primary total knee replacement at a minimum of six months after operation. The knee replacement used in all the patients was an unconstrained, uncemented, hydroxyapatite-coated posterior-cruciate-ligament-retaining prosthesis with the option of a cemented polyethylene patellar button. All patients had a medial parapatellar skin incision. We recorded age, the time from surgery, the Knee Society score (KSS),⁸ the range of movement and whether a patellar button was implanted at arthroplasty.

The patients were asked about their ability to kneel and then to demonstrate kneeling on a firm surface and to record the level of pain experienced on a scale of from 0 to 10. Those patients unwilling or unable to kneel were asked to explain the reason (knee pain, back or hip stiffness, anxiety, etc) and this was recorded. Thus two groups emerged: those able to kneel without pain or with only mild pain (score 0 to 4) and those unable to kneel because of pain in the knee (5 to 10). Those patients who stated that they were unable to kneel because of back pain, stiffness of the hip, etc, were excluded from this group.

The mean age of the patients was 66 years (43 to 82) with a mean time from surgery of 30 months (6 to 127). The mean knee score was 179/200 (123 to 200). The mean range of movement was 1° to 114° (-5 to 135). The number of knees with patellar resurfacing was 35.

S. H. Palmer, FRCS (Trauma & Orth), Locum Consultant Orthopaedic Surgeon
Nuffield Orthopaedic Centre, Windmill Road, Headington, Oxford OX3 74D, UK.

C. T. Servant, FRCS, Knee Fellow
J. Maguire, FRACS Ortho, Knee Fellow
E. N. Parish, BHSc, HM, Research Assistant
M. J. Cross, OAM, MD, FRACS, Consultant Orthopaedic Surgeon
The Australian Institute of Musculo-Skeletal Research, 286 Pacific Highway, Crows Nest, New South Wales, Australia 2065.

Correspondence should be sent to Mr S. H. Palmer at 3 Mark Road, Headington, Oxford OX3 8PB, UK.

©2002 British Editorial Society of Bone and Joint Surgery
0301-620X/02/212568 \$2.00

We compared and analysed differences in patellar resurfacing, range of movement and knee scores between those who were able to kneel and those who could not using SPSS 10.0 (SPSS Inc, Chicago, Illinois) statistical software (Fisher's exact test, Mann-Whitney U test).

Results

In 32 knees patients stated that they could kneel without significant discomfort. In 54 knees patients avoided kneeling because of uncertainties or recommendations from third parties (doctors, nursing staff, friends, etc). A total of 64 patients was actually able to kneel without pain or discomfort or with mild discomfort only. Twelve were unable to kneel because of problems which were not related to the knee. Twenty-four were therefore unable to kneel because of discomfort in the knee.

Of the knees with patellar resurfacing, 26 were in patients who were able to kneel and five in those who could not. Of those without resurfacing, 38 were in patients who were able to kneel and 19 in those who could not ($p = 0.1$).

The mean range of movement was 114° in the knees of patients who were able to kneel and 110° in those who could not. The mean KSS was 183 in the knees of patients who were able to kneel and 179 in those who could not. These differences were not significant for both range of movement ($p = 0.6$) and KSS ($p = 0.6$).

Discussion

Our results suggest that there are uncertainties about the ability to kneel after total knee replacement. In 32 knees patients declared that they had the ability to kneel although in 64 knees they were actually able to demonstrate the ability to kneel comfortably. Of those patients who did not kneel before the study 80% avoided kneeling because of advice from medical staff or third parties or because of fear of harming the prosthesis. The ability to kneel did not appear to relate to whether or not the patella had been resurfaced, the range of movement of the knee or the KSS.

Radiographs showed two patterns of kneeling (Figs 1 and 2). 'Upright kneeling' which occurs with 90° of knee flexion and 'flexed kneeling' which can occur when more than 110° of knee flexion is achieved. In the first the points of contact with the ground are the patella and the tibial tuberosity. In the second, only the tibial tuberosity bears weight. In our study the patients did not localise the source of discomfort when kneeling, but it would seem that kneeling in a flexed position ($>110^\circ$) reduces the forces across the patellofemoral articulation.

Negative advice given to patients regarding kneeling seems to have arisen out of concern for the prosthesis, but there are no published data concerning this risk. Examination of the radiographs taken when kneeling suggests that



Fig. 1

Lateral radiograph of a knee in a patient in an 'upright' kneeling position.




Fig. 2

Lateral radiograph of the knee in a patient in the 'flexed' kneeling position.



Fig. 3

Lateral radiograph showing upright kneeling in a normal knee.

compressive forces through the femoral component are similar to those experienced in normal standing and walking (Figs 1 and 2). The tibial tray appears to be protected since it has no contact with the weight-bearing surface. The patellar button, when present, is in compression and, because of its obliquity, appears to experience a degree of shear. This differs from the normal knee (Fig. 3). Kneeling tends to produce posterior subluxation of the tibia, but this is prevented by the PCL and the congruency of the bearing surface and the tibial post in a PCL-retained replacement system. Therefore the forces through the prosthesis appear theoretically to be benign except possibly for the patellar button and the tibial tray in PCL-sacrificing devices.

The only previous publication on this subject showed similar results. Schai, Gibbon and Scott¹⁶ found that 44% of their patients stated that they could kneel, but 82% were actually observed to kneel easily. Fear of harming the prosthesis and lack of information explained why 49% did not kneel. Differences in operative technique from our study included a midline skin incision and patellar resurfacing in 89% of patients. Both studies recorded differences between the perceived and actual ability to kneel. Minor differences between the design of the studies may account for the different figures. Patients with moderate or severe pain on kneeling were categorised as 'non-kneelers' in our study.

The use of a lateral parapatellar skin incision has been shown to produce fewer neuromata and kneeling difficulties after surgical exploration of ligamentous injuries.¹⁷ We prefer to use a medial parapatellar skin incision in total knee arthroplasty to prevent elevation of potentially devitalised skin flaps at the front of the knee.¹⁸ The risk of poor wound healing has to be weighed against the marginally improved ability to kneel with a lateral skin incision.

The actual ability to kneel after total knee replacement differs from the perceived ability to kneel and varies between 64% and 82% on current evidence. Fear of harming the prosthesis or uncertainties about recommendations on kneeling are the main reasons for this difference. It does not appear to be related to the presence of patellar resurfacing, the range of movement or the functional knee score.

No benefits in any form have been received or will be received from a commercial party related directly or indirectly to the subject of this article.

References

1. Rorabeck CH, Bourne RB, Lewis PL, Nott L. The Miller-Galante knee prosthesis for the treatment of osteoarthritis: a comparison of the results of partial fixation with cement and fixation without any cement. *J Bone Joint Surg [Am]* 1993;75-A:402-8.
2. Schai PA, Thornhill TS, Scott RD. Total knee arthroplasty with the PFC system: results at a minimum of ten years and survivorship analysis. *J Bone Joint Surg [Br]* 1998;80-B:850-8.
3. Robertsson O, Scott G, Freeman MAR. Ten-year survival of the cemented Freeman-Samuels primary knee arthroplasty: data from the Swedish knee arthroplasty register and the Royal London Hospital. *J Bone Joint Surg [Br]* 2000;82-B:506-7.
4. Wright J, Ewald FC, Walker PS, et al. Total knee arthroplasty with the kinematic prosthesis: results after five to nine years: a follow-up note. *J Bone Joint Surg [Am]* 1990;72-A:1003-9.
5. Mokris JG, Smith SW, Anderson SE. Primary total knee arthroplasty using the Genesis total knee arthroplasty system: 3- to 6-year follow-up study of 105 knees. *J Arthroplasty* 1997;12:91-8.
6. Knutson K, Lewold S, Robertsson O, Lidgren L. The Swedish knee arthroplasty register: a nationwide study of 30,003 knees 1976-1992. *Acta Orthop Scand* 1994;65:375-86.
7. Insall JN, Ranawat CS, Aglietti P, Shine J. A comparison of four models of total knee-replacement prostheses. *J Bone Joint Surg [Am]* 1976;58-A:754-65.
8. Insall JN, Dorr LD, Scott RD, Scott WN. Rationale of the Knee Society clinical rating system. *Clin Orthop* 1989;248:13-4.
9. Lequesne MG, Mery C, Samson M, et al. Indexes of severity for osteoarthritis of the hip and knee: validation - value in comparison with other assessment tests. *Scand J Rheumatol Suppl* 1987;65:85-9.
10. Irrgang JJ, Snyder-Mackler L, Wainner RS, Fu FN, Harner CD. Development of a patient-reported measure of function of the knee. *J Bone Joint Surg [Am]* 1998;80-A:1132-45.
11. Bellamy N, Buchanan WW, Goldsmith CH, Campbell J, Stitt LW. Validation study of WOMAC: a health status instrument for measuring clinically important patient relevant outcomes to antirheumatic drug therapy in patients with osteoarthritis of the hip or knee. *J Rheumatol* 1988;15:1833-40.
12. Dawson J, Fitzpatrick R, Murray D, Carr A. Questionnaire on the perceptions of patients about total knee replacement. *J Bone Joint Surg [Br]* 1998;80-B:63-9.
13. Cooper C, McAlindon T, Coggon D, Egger P, Dieppe P. Occupational activity and osteoarthritis of the knee. *Ann Rheum Dis* 1994;53:90-3.
14. Coggon D, Croft P, Kellingray S, et al. Occupational physical activities and osteoarthritis of the knee. *Arthritis Rheum* 2000;43:1443-9.
15. Ulbrich J, Raheja A, Alexander NB. Body positions used by healthy and frail older adults to rise from the floor. *J Am Geriatr Soc* 2000;48:1626-32.
16. Schai PA, Gibbon AJ, Scott RD. Kneeling ability after total knee arthroplasty: perception and reality. *Clin Orthop* 1999;367:195-200.
17. Berg P, Mjoberg B. A lateral skin incision reduces peripatellar dysesthesia after knee surgery. *J Bone Joint Surg [Br]* 1991;73-B:374-6.
18. Klein NE, Cox CV. Wound problems in total knee arthroplasty. In: Fu FH, Harner CD and Vince KG, eds. *Knee surgery* Baltimore: Williams and Wilkins, 1994:1539-52.